



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

divisions in the ascus follow the same routine as described by Miss FRASER in *Humaria rutilans*. The first mitosis is heterotypic, the second homotypic, and the third brachymeiotic, leading to a reduction without division in the last mitosis. The spores are formed by the astral rays in the manner described by HARPER for *Pyronema*.—H. HASSELBRING.

**A parasitic orchid.**—According to the investigation of KUSANO,<sup>46</sup> the vegetative body of the orchid *Gastrodia elata* consists of a tuberous rhizome which multiplies through the production of tuberous offshoots. If these offshoots form no mycorrhiza, they decrease in size and finally die without being able to flower, but if infected by the mycelium of *Armillaria mellea* they enlarge, flower, and produce daughter tubercles; it is therefore concluded that the orchid is completely parasitic upon the fungus. The mycorrhiza is of the ectotropic type; occasionally, however, the fungus behaves as a parasite, penetrating deeply into the tissue of the tubers and causing their collapse in a manner similar to that seen in potato tubers attacked by the same organism.—GEO. D. FULLER.

**Budding in Cycas.**—While in Japan, Miss STOPES<sup>47</sup> noted the origin of the well known adventitious buds of *Cycas revoluta*. They arise on the upper portions of old leaf bases, some appearing more than 200 crowns back of the growing point. None were found arising from the axis, and the young buds do not seem to have any connection with the axis. The buds when removed grow into normal plants, but when they develop strongly upon the parent plant they give rise to the so-called "branching," and in such cases, of course, vascular connections are established with the main axis.—CHARLES J. CHAMBERLAIN.

---

<sup>46</sup> KUSANO, S., Preliminary note on *Gastrodia elata* and its mycorrhiza. Ann. Botany 25:521-523. 1911.

<sup>47</sup> STOPES, MARIE C., Adventitious budding and branching in *Cycas revoluta*. New Phytol. 9:234-241. figs. 7. 1910.